

Correlations between lepton flavour violation at the LHC and type-I seesaw parameters in minimal supergravity

A. Villanova del Moral*

Departamento de Física and CFTP, Instituto Superior Técnico, Avenida Rovisco Pais 1, 1049-001 Lisboa, Portugal

The most general supersymmetric seesaw mechanism has too many parameters to be predictive and thus can not be excluded by *any* measurements of lepton flavour violating (LFV) processes. We focus on the simplest version of the type-I seesaw mechanism assuming minimal supergravity (mSugra) boundary conditions. We compute branching ratios for the LFV scalar tau decays, $\tilde{\tau}_2 \rightarrow (e, \mu) + \chi_1^0$, as well as loop-induced LFV decays at low energy, such as $l_i \rightarrow l_j + \gamma$ and $l_i \rightarrow 3l_j$, exploring their sensitivity to the unknown seesaw parameters. We find some simple, extreme scenarios for the unknown right-handed parameters, where ratios of LFV branching ratios correlate with neutrino oscillation parameters. If the overall mass scale of the left neutrinos and the value of the reactor angle were known, the study of LFV allows, in principle, to extract information about the so far unknown right-handed neutrino parameters.

1. INTRODUCTION

Neutrino experiments [1] have firmly established that neutrinos are massive and global fits to all neutrino oscillation data [2] give precise values for the neutrino mass splittings and their mixing angles. The evidence of massive neutrinos provides the first experimental signal of physics beyond the Standard Model, being the so-called seesaw mechanism [3] the most popular mechanism to generate Majorana neutrino masses.

In this work [4] we study the correlation with neutrino parameters of the ratio of LFV decays of the stau, assuming mSugra boundary conditions and the simplest type-I seesaw mechanism as the origin of neutrino masses and mixings. In this theoretical framework, left-slepton LFV decays are proportional to the square of the off-diagonal elements of the slepton mass matrix, due to the renormalization group equation (RGE) running of the soft breaking parameters.

2. ANALYSIS

For qualitative understanding, we consider the leading-log approximated solutions to the RGE's. The left-slepton LFV decays are then proportional to

$$\text{Br}(\tilde{l}_i \rightarrow l_j \chi_1^0) \propto |(\Delta M_{\tilde{L}}^2)_{ij}|^2 \propto |(Y_\nu^\dagger L Y_\nu)_{ij}|^2. \quad (1)$$

We can parametrize the neutrino Yukawa matrix as [6]

$$Y_\nu = \sqrt{2} \frac{i}{v_U} \sqrt{\hat{M}_R} R \sqrt{\hat{m}_\nu} U^\dagger, \quad (2)$$

where $\hat{m}_\nu$ and $\hat{M}_R$ are the diagonal matrices with the light neutrino mass eigenvalues m_i and the heavy neutrino mass eigenvalues M_i , respectively; U is the leptonic mixing matrix and R is a complex orthogonal matrix. This way, the left-slepton LFV decays are correlated to neutrino parameters

$$\text{Br}(\tilde{l}_i \rightarrow l_j \chi_1^0) \propto \left| U_{i\alpha} U_{j\beta}^* \sqrt{m_\alpha} \sqrt{m_\beta} R_{k\alpha}^* R_{k\beta} M_k \log \left(\frac{M_X}{M_k} \right) \right|^2. \quad (3)$$

*Electronic address: albert@cftp.ist.utl.pt

In order to eliminate most of the dependence on the supersymmetric parameters, we work with ratios of LFV decays. Thus, for example, the ratio of stau LFV decays can be expressed in terms of the parameter r_{23}^{13} ,

$$\frac{\text{Br}(\tilde{\tau}_2 \rightarrow e + \chi_1^0)}{\text{Br}(\tilde{\tau}_2 \rightarrow \mu + \chi_1^0)} \simeq \frac{|(\Delta M_L^2)_{13}|^2}{|(\Delta M_L^2)_{23}|^2} \equiv (r_{23}^{13})^2, \quad (4)$$

which only depends on neutrino parameters.

In the case of degenerate right-handed neutrinos and assuming R being real and tribimaximal (TBM) mixing [5],

$$r_{23}^{13} = \frac{2(m_2 - m_1)}{|3m_3 - 2m_2 - m_1|}. \quad (5)$$

Table I shows the form of Eq. (5) and its numerical values, for different neutrino scenarios. For the case $s_{13} \neq 0$,

Table I: Parameter r_{23}^{13} for the case of degenerate right-handed neutrinos, assuming R being real and TBM mixing. Each column corresponds to a different neutrino scenario: strict normal hierarchical (SNH), strict inverse hierarchical (SIH), quasidegenerate normal hierarchical (QDNH) and quasidegenerate inverse hierarchical (QDIH) neutrinos. First row shows the analytical form of the parameter r_{23}^{13} ; second row (BFP) shows its value when fixing neutrino mass splittings to their best fit point value and third row shows its value when considering the 3σ allowed range for the neutrino mass-squared differences. Note that $\alpha \equiv \Delta m_s^2/|\Delta m_\lambda^2|$ is the ratio of the solar over the atmospheric mass splitting and $\sigma_\lambda \equiv \Delta m_\lambda^2/|\Delta m_\lambda^2|$ is the sign of the atmospheric mass splitting.

	SNH	SIH	QDNH	QDIH
	$r_{23}^{13} = \frac{2\sqrt{\alpha}}{3\sqrt{1+\alpha}-2\sqrt{\alpha}}$	$r_{23}^{13} = \frac{2(1-\sqrt{1-\alpha})}{2+\sqrt{1-\alpha}}$	$r_{23}^{13} \simeq \frac{2\alpha}{3\sigma_\lambda+\alpha}$	
BFP	$(r_{23}^{13})^2 = 1.7 \times 10^{-2}$	$(r_{23}^{13})^2 = 1.1 \times 10^{-4}$	$(r_{23}^{13})^2 = 4.4 \times 10^{-4}$	$(r_{23}^{13})^2 = 4.6 \times 10^{-4}$
3σ	$(r_{23}^{13})^2 \in [0.92, 3.7] \times 10^{-2}$	$(r_{23}^{13})^2 \in [0.48, 3.3] \times 10^{-4}$	$(r_{23}^{13})^2 \in [1.8, 12] \times 10^{-4}$	$(r_{23}^{13})^2 \in [1.9, 13] \times 10^{-4}$

Fig. 1 shows the squared ratios as a function of s_{13}^2 , for different neutrino scenarios and for two choices of the Dirac phase δ . For strongly hierarchical right-handed neutrinos and assuming R being the identity, the parameter r_{23}^{13} does




Figure 1: Square ratios $(r_{13}^{13})^2$ (blue line, dotted line), $(r_{23}^{13})^2$ (red line, dashed line) and $(r_{23}^{13})^2$ (green line, full line) versus s_{13}^2 for the case of degenerate heavy neutrinos, real R and the rest of the neutrino parameters fixed to their best fit point values. Each column correspond to a different neutrino scenario: SNH (first row), SIH (second row), QDNH (third row) and QDIH (forth row). Each row correspond to a different value of the Dirac phase: $\delta = 0$ (first column) and $\delta = \pi$ (second column).

not depend on the light neutrino masses any more, but on their mixing angles and Dirac phase. For the specific case in which $s_{13} = 0$, Table II shows the analytical form of r_{23}^{13} and its numerical values, for different right-handed neutrino dominant mass scenarios. For the case of nonzero values of s_{13} , Fig. 2 shows the squared ratios as a function

Table II: Parameter r_{23}^{13} for the case of strongly hierarchical right-handed neutrinos, assuming R being the identity and $s_{13} = 0$. Each column corresponds to a different heavy neutrino scenario: if M_1 is the heaviest mass eigenvalue (Dominant M_1), if M_2 is the heaviest mass eigenvalue (Dominant M_2) and if M_3 is the heaviest mass eigenvalue (Dominant M_3). First row shows the analytical form of the parameter r_{23}^{13} ; second row (BFP) shows its value when fixing neutrino mixing angles to their best fit point value and third row shows its value when considering the 3σ allowed range for the neutrino mixing angles.

	Dominant M_1	Dominant M_2	Dominant M_3
	$r_{23}^{13} = \frac{c_{12}}{s_{12}c_{23}}$	$r_{23}^{13} = \frac{s_{12}}{c_{12}c_{23}}$	$r_{23}^{13} = 0$
BFP	$(r_{23}^{13})^2 = 4$	$(r_{23}^{13})^2 = 1$	$(r_{23}^{13})^2 = 0$
3σ	$(r_{23}^{13})^2 \in [2.3, 8.6]$	$(r_{23}^{13})^2 \in [0.53, 2.0]$	$(r_{23}^{13})^2 = 0$

of s_{13}^2 , for different right-handed neutrino dominant mass scenarios. Note that for the scenario of dominant M_3 , the parameter r_{23}^{13} is proportional to s_{13} .




Figure 2: Square ratios $(r_{13}^{12})^2$ (blue line, dotted line), $(r_{23}^{12})^2$ (red line, dashed line) and $(r_{23}^{13})^2$ (green line, full line) versus s_{13}^2 for the case of strongly hierarchical heavy neutrinos, R being the identity and the rest of the neutrino parameters fixed to their best fit point values. First and second panel correspond to $\delta = 0$ and $\delta = \pi$, respectively, in the case of dominant M_1 , while third and fourth panel correspond to $\delta = 0$ and $\delta = \pi$, respectively, in the case of dominant M_2 .

3. RESULTS

In order to check the validity of our analytical estimated ratio of stau LFV decays, we have performed a numerical calculation of such LFV decays making use of the program package SPHENO [7]. More details can be found in Ref. [4].

For degenerate right-handed neutrinos, first panel in Fig. 3 shows the stau LFV decays as a function of the heavy neutrino common mass M_R , for mSugra benchmark point SPS3 [8], SNH ($m_1 = 0$) and TBM mixing.

For strongly hierarchical right-handed neutrinos, the other panels in Fig. 3 show the stau LFV decays as a function of the heavy neutrino dominant mass, which is M_1 in the second panel, M_2 in the third panel and M_3 in the fourth panel. Again, NH and TBM have been assumed.

Note that the ratio of the stau LFV decays follows very accurately the analytical estimate in the region allowed by the upper limit on $\text{Br}(\mu \rightarrow e\gamma)$.

4. CONCLUSION

Neutrino experimental data show that neutrinos are massive and mix. If its origin is the simplest supersymmetric type-I seesaw mechanism and mSugra boundary conditions hold, then LFV decays are related to neutrino parameters. In particular, we have studied the relation of the ratio of the stau LFV decays with neutrino parameters for different neutrino scenarios.




Figure 3: Left-stau LFV decays as a function of the right-handed neutrino degenerate common mass M_R (first panel), of the dominant M_1 (second panel), of the dominant M_2 (third panel) and of the dominant M_3 (fourth panel), for the SPS3 standard point, R being the identity, TBM mixing and SNH (except for the second panel, where $m_1 = 0.1$ eV). In the panels where one M_i is dominant, the other two heavy neutrino masses have been fixed to 10^{10} GeV.

Acknowledgments

The author wish to thank his collaborators M. Hirsch, W. Porod, J. C. Romao and J. W. F. Valle. The author is supported by *Fundação para a Ciência e a Tecnologia* under the grant SFRH/BPD/30450/2006. This work was partially supported by *Fundação para a Ciência e a Tecnologia* through the projects POCI/81919/2007 and CFTP-FCT UNIT 777, which are partially funded through POCTI (FEDER) and by the Marie Curie RTNs MRTN-CT-2006-035505 and MRTN-CT-2004-503369. It was also partially supported by Spanish grants FPA2005-01269 and Accion Integrada HA-2007-0090 (MEC) and by the European Commission network MRTN-CT-2004-503369 and ILIAS/N6 RII3-CT-2004-506222.

References

- [1] Y. Fukuda *et al.* [Super-Kamiokande Collaboration], Phys. Rev. Lett. **81**, 1562 (1998) [arXiv:hep-ex/9807003]. S. Abe *et al.* [KamLAND Collaboration], Phys. Rev. Lett. **100**, 221803 (2008) [arXiv:0801.4589 [hep-ex]]. C. Arpesella *et al.* [Borexino Collaboration], arXiv:0805.3843 [astro-ph]. B. Aharmim *et al.* [SNO Collaboration], Phys. Rev. Lett. **101**, 111301 (2008) [arXiv:0806.0989 [nucl-ex]]. P. Adamson *et al.* [MINOS Collaboration], arXiv:0806.2237 [hep-ex].
- [2] M. Maltoni, T. Schwetz, M. A. Tortola and J. W. F. Valle, New J. Phys. **6**, 122 (2004) [arXiv:hep-ph/0405172]; The arXiv version 6 provides results updated as of September 2007. T. Schwetz, M. Tortola and J. W. F. Valle, arXiv:0808.2016 [hep-ph].
- [3] P. Minkowski, Phys. Lett. B **67**, 421 (1977). M. Gell-Mann, P. Ramond and R. Slansky, in *Supergravity*, edited by P. van Nieuwenhuizen and D. Freedman (North Holland, Amsterdam, 1979). T. Yanagida, in *Proceedings of the Workshop on the Baryon Number of the Universe and Unified Theories*, edited by O. Sawada and A. Sugamoto (Tsukuba, Japan, 1979). R. N. Mohapatra and G. Senjanovic, Phys. Rev. Lett. **44**, 912 (1980). J. Schechter and J. W. F. Valle, Phys. Rev. D **22**, 2227 (1980). J. Schechter and J. W. F. Valle, Phys. Rev. D **25**, 774 (1982).
- [4] M. Hirsch, J. W. F. Valle, W. Porod, J. C. Romao and A. Villanova del Moral, Phys. Rev. D **78**, 013006 (2008) [arXiv:0804.4072 [hep-ph]].
- [5] P. F. Harrison, D. H. Perkins and W. G. Scott, Phys. Lett. B **530**, 167 (2002) [arXiv:hep-ph/0202074].
- [6] J. A. Casas and A. Ibarra, Nucl. Phys. B **618**, 171 (2001) [arXiv:hep-ph/0103065].
- [7] W. Porod, Comput. Phys. Commun. **153**, 275 (2003) [arXiv:hep-ph/0301101].
- [8] B. C. Allanach *et al.*, Eur. Phys. J. C **25**, 113 (2002) [arXiv:hep-ph/0202233]. J. A. Aguilar-Saavedra *et al.*, Eur. Phys. J. **C46**, 43 (2006) [arXiv:hep-ph/0511344].